arXiv:1612.04450v1 [math.AT] 14 Dec 2016

STRICT n-CATEGORIES AND AUGMENTED DIRECTED COMPLEXES MODEL HOMOTOPY TYPES

ANDREA GAGNA

Abstract. In this paper we show that both the homotopy category of strict n-categories, $1 \leqslant n \leqslant \infty$, and the homotopy category of Steiner's augmented directed complexes are equivalent to the category of homotopy types. In order to do so, we first prove an abstract result, based on a strategy of Fritsch and Latch, giving sufficient conditions for a nerve functor with values in simplicial sets to induce an equivalence at the level of homotopy categories. We then apply this result to strict n-categories and augmented directed complexes, for which the hypothesis of our theorem were established by Ara and Maltsiniotis.

Introduction

The homotopy theory of small categories was born with the introduction by Grothendieck of the nerve functor in [5], allowing the definition of weak equivalences in $\mathcal{C}at$, the category of small categories: a functor is a weak equivalence precisely when its nerve is a simplicial weak equivalence. This definition provides us with an induced nerve functor at the level of the homotopy categories, that is the categories in which we localise at weak equivalences.

The first striking result of this theory appears in Illusie's thesis [7] (who credits it to Quillen) and it states that the induced nerve functor is an equivalence of homotopy categories. A careful study of the subtle homotopy theory of small categories by Thomason [11], lead him to show another important result: the existence of a model structure on $\mathcal{C}at$ which is Quillen equivalent to the Kan-Quillen model structure on simplicial sets. This implies that small categories and simplicial sets are not only equivalent as homotopy categories, but actually as $(\infty, 1)$-categories.

Similarly Chiche [3], inspired by ideas of del Hoyo [6], has showed that strict 2-categories model homotopy types. Moreover, Ara and Maltsiniotis [2] provided an abstract framework in which a Quillen equivalence between n-$\mathcal{C}at$ and simplicial sets can be established, using the Street nerve, as long as some conditions are satisfied; they also showed that these conditions hold for small 2-categories. In a subsequent paper [1] they proved in full generality one of these conditions, namely condition (e), making an important use of augmented directed complexes introduced by Steiner [9], which act as a linearisation of ∞-categories and provide a meaningful nerve functor to simplicial sets.

In the same article, they also conjectured that for every n such that $1 \leqslant n \leqslant \infty$, the Street nerve induces an equivalence of homotopy categories between n-$\mathcal{C}at$ and simplicial sets and that the same holds for augmented directed complexes with their own nerve functor. The aim of this short paper is to prove these conjectures.

We consider a general framework for a nerve $U\colon \mathcal{C} \to \widehat{\mathbf{\Delta}}$ induced as a right adjoint of a left Kan extension F of a cosimplicial object of $\mathcal{C}$ along the Yoneda

embedding of $\mathbf{\Delta}$, where $\mathcal{C}$ is a cocomplete category. Using the simplicial properties of subdivision and adapting a strategy developed in an article by Fritsch and Latch [4], a short and abstract proof that U induces an equivalence at the level of homotopy categories can be given, as long as it satisfies an appropriate version of condition (e). This condition says that the unit $\eta\colon 1_{\widehat{\mathbf{\Delta}}} \to UF$ is a weak equivalence on the nerves of posets, that is for every poset E, the morphism of simplicial sets $N_1(E) \to UF(N_1(E))$ is a weak equivalence, where $N_1\colon \mathcal{C}at \to \widehat{\mathbf{\Delta}}$ denotes the usual nerve. Ara and Maltsiniotis have established this condition for both the nerve $N_m\colon m\text{-}\mathcal{C}at \to \widehat{\mathbf{\Delta}}$ of strict m-categories, $1 \leqslant n \leqslant \infty$, and the nerve $N\colon \mathcal{C}_{\mathrm{da}} \to \widehat{\mathbf{\Delta}}$ of augmented directed complexes. We can thus apply our theorem to these cases and get a proof of the above conjectures.

Let us fix some notations and terminology. If A is a small category, the category of presheaves on A will be denoted by $\widehat{A}$. If F and G are two functors between two categories $\mathcal{C}$ to $\mathcal{D}$ and W is a class of arrows of $\mathcal{D}$ called weak equivalences, we will say that a natural transformation $\alpha\colon F \to G$ is a *weak equivalence* when it is a component-wise weak equivalence, i.e., for any object c of $\mathcal{C}$ the arrow α_c is an element of W. For any adjunction, we will call η the unit and ε the counit.

1. Simplicial preliminaries

1.1. The *category of simplices* $\mathbf{\Delta}$ is the category whose objects are the finite ordinals $\Delta_n = \{0, 1, \dots, n\}$, for $n \geqslant 0$, and whose arrows are the order-preserving maps; the *category of semi-simplices* $\mathbf{\Delta}'$ is the category with the same objects as $\mathbf{\Delta}$ but only strictly increasing maps. We have a canonical inclusion functor $i\colon \mathbf{\Delta}' \to \mathbf{\Delta}$. We will denote by $i^*\colon \widehat{\mathbf{\Delta}} \to \widehat{\mathbf{\Delta}'}$ the functor given by precomposition by i and by $i_!\colon \widehat{\mathbf{\Delta}'} \to \widehat{\mathbf{\Delta}}$ its left adjoint.

We denote by $\mathcal{C}at$ the category of small categories and by $N_1\colon \mathcal{C}at \to \widehat{\mathbf{\Delta}}$ and $c_1\colon \widehat{\mathbf{\Delta}} \to \mathcal{C}at$ the usual nerve and its left adjoint.

1.2. We shall call *weak equivalences of* $\widehat{\mathbf{\Delta}}$ the usual weak homotopy equivalences and *weak equivalences of* $\widehat{\mathbf{\Delta}'}$ the morphisms such that their image by $i_!$ is a weak equivalence of $\widehat{\mathbf{\Delta}}$.

Proposition 1.3. *The adjunction morphisms* $\eta\colon 1_{\widehat{\mathbf{\Delta}'}} \to i^*i_!$ *and* $\varepsilon\colon i_!i^* \to 1_{\widehat{\mathbf{\Delta}}}$ *are weak equivalences.*

Proof. See Proposition 2.12 of [4]. □

Corollary 1.4. *The functors* $i_!$ *and* i^* *both respect weak equivalences and induce an equivalence pair at the level of homotopy categories.*

Proof. This follows immediately from the previous proposition. □

1.5. We denote by $\mathrm{Sd}\colon \widehat{\mathbf{\Delta}} \to \widehat{\mathbf{\Delta}}$ the subdivision endofunctor of simplicial sets introduced by Kan in [8, §7] and by $\alpha\colon \mathrm{Sd} \to 1_{\widehat{\mathbf{\Delta}}}$ the canonical natural transformation. Recall that α is a weak equivalence (see Lemma 7.5 of loc. cit.). Therefore the previous proposition implies that the natural transformation

$$\alpha^2 * \varepsilon\colon \mathrm{Sd}^2 i_! i^* \to 1_{\widehat{\mathbf{\Delta}}}$$

is a weak equivalence, where $\alpha^2 = \alpha * \alpha$.

Proposition 1.6. *For any simplicial set* X*, the simplicial set* $\mathrm{Sd}^2 i_! i^*(X)$ *is isomorphic to the nerve of a poset.*

Proof. By Corollary 3.11 and Remark 3.15 of [4], the simplicial set $\mathrm{Sd}^2 i_! i^*(X)$ is isomorphic to the nerve of a small category, say $N_1(A)$. It is in fact isomorphic to the nerve of the category $c_1\mathrm{Sd}^2 i_! i^*(X)$, as we have

$$\mathrm{Sd}^2 i_! i^*(X) \cong N_1(A) \cong N_1 c_1 N_1(A) \cong N_1 c_1 \mathrm{Sd}^2 i_! i^*(X).$$

But, for any simplicial set Y, the category $c_1\mathrm{Sd}^2(Y)$ is a poset (see Lemma 5.6 of [11]). □

2. An abstract Illusie-Quillen theorem

2.1. We fix $\mathcal{C}$ a cocomplete category and $j\colon \mathbf{\Delta} \to \mathcal{C}$ a cosimplicial object in $\mathcal{C}$. We denote by U the induced nerve functor, right adjoint to $F\colon \widehat{\mathbf{\Delta}} \to \mathcal{C}$ the left Kan extension of j along the Yoneda embedding of $\mathbf{\Delta}$. We call *weak equivalences of* $\mathcal{C}$ the morphisms sent by U to weak equivalences of $\widehat{\mathbf{\Delta}}$. In particular, U induces a functor at the level of the homotopy categories.

2.2. We shall say that j *satisfies condition (e)* if the unit $\eta\colon 1_{\widehat{\mathbf{\Delta}}} \to UF$ is a weak equivalence on the nerves of posets, i.e., if the morphism $N_1(E) \to UF(N_1(E))$ of simplicial sets is a weak equivalence for any poset E.

Remark 2.3. We named this property after the "condition (e)" appearing in the abstract Thomason Theorem (Theorem 4.11 of [2]) of Ara and Maltsiniotis.

Theorem 2.4 (Abstract Illusie–Quillen Theorem)**.** *Let $j\colon \mathbf{\Delta} \to \mathcal{C}$ be a cosimplicial object satisfying condition (e). Then the functor $U\colon \mathcal{C} \to \widehat{\mathbf{\Delta}}$ induces an equivalence at the level of the homotopy categories.*

Proof. Let $Q\colon \widehat{\mathbf{\Delta}} \to \widehat{\mathbf{\Delta}}$ be a functor such that for every simplicial set X, the simplicial set $Q(X)$ is isomorphic to the nerve of a poset and $\gamma\colon Q \to 1_{\widehat{\mathbf{\Delta}}}$ a natural transformation which is a weak equivalence. For instance, we can choose the functor $\mathrm{Sd}^2 i_! i^*$ as Q by virtue of Proposition 1.6 and as γ the natural transformation $\mathrm{Sd}^2 i_! i^* \to 1_{\widehat{\mathbf{\Delta}}}$ considered in paragraph 1.5.

We will show that FQ respects weak equivalences and gives an inverse in homotopy of the functor induced by U. We have a zig-zag of natural transformations

$$UFQ \xleftarrow{\eta * Q} Q \xrightarrow{\gamma} 1_{\widehat{\mathbf{\Delta}}}\ .$$

Condition (e) and the fact that Q takes values in nerves of posets show that $\eta * Q$ is a weak equivalence; by definition, γ is so, too. This establishes that FQ preserves weak equivalences and that it is a left inverse for U in homotopy.

Moreover, we have a natural transformation

$$FQU \xrightarrow{F*\gamma*U} FU \xrightarrow{\varepsilon} 1_{\mathcal{C}}\ .$$

We have to show that

$$UFQU \xrightarrow{UF*\gamma*U} UFU \xrightarrow{U*\varepsilon} U$$

is a weak equivalence. Considering the commutative diagram



we note that $\eta * QU$ is a weak equivalence for the same reason as above, that is by condition (e), and that $\gamma * U$ is a weak equivalence by definition. By virtue of the 2-out-of-3 for the weak equivalences of $\mathcal{C}$, this proves the theorem. □

3. Strict n-categories and augmented directed complexes model homotopy types

3.1. For any n such that $1 \leqslant n \leqslant \infty$, we denote by n-$\mathcal{C}at$ the category of small strict n-categories. We recall that for any positive integer m, the full inclusion m-$\mathcal{C}at \to \infty$-$\mathcal{C}at$ has a left adjoint τ_m^i.

3.2. Street introduced in [10] a cosimplicial object $\mathcal{O}\colon \mathbf{\Delta} \to \infty$-$\mathcal{C}at$, from which we can define a pair of adjoint functors

$$c_\infty\colon \widehat{\mathbf{\Delta}} \to \infty\text{-}\mathcal{C}at \quad \text{and} \quad N_\infty\colon \infty\text{-}\mathcal{C}at \to \widehat{\mathbf{\Delta}}.$$

The right adjoint N_∞ is known as the *Street nerve*. For any positive integer m, we can define a functor $\mathcal{O}^{\leqslant m} = \tau_m\mathcal{O}\colon \mathbf{\Delta} \to m$-$\mathcal{C}at$ and therefore a pair of adjoint functors

$$c_m\colon \widehat{\mathbf{\Delta}} \to \infty\text{-}\mathcal{C}at \quad \text{and} \quad N_m\colon \infty\text{-}\mathcal{C}at \to \widehat{\mathbf{\Delta}}.$$

The functor N_m is the restriction of the Street nerve to the full subcategory m-$\mathcal{C}at$ and c_m is the composition of the realisation functor c_∞ followed by τ_m^i. When $m = 1$, we get the usual nerve functor.

3.3. For any m, $1 \leqslant m \leqslant \infty$, we define the *weak equivalences of* m-$\mathcal{C}at$ as the morphisms whose image by N_m is a weak equivalence of $\widehat{\mathbf{\Delta}}$. These are also called *Thomason weak equivalences* in [2].

Theorem 3.4. *For any m such that $1 \leqslant m \leqslant \infty$, the functor $N_m\colon m$-$\mathcal{C}at \to \widehat{\mathbf{\Delta}}$ induces an equivalence at the level of the homotopy categories. In particular, strict n-categories model homotopy types.*

Proof. By Theorem 2.4, it is enough to show that the cosimplicial objects $\mathcal{O}^{\leqslant m}$ from $\mathbf{\Delta}$ to m-$\mathcal{C}at$ satisfy condition (e) for any m, $1 \leqslant m \leqslant \infty$. This is precisely Corollary 10.11 of [1]. □

3.5. We shall denote by $\mathcal{C}_{\mathrm{da}}$ the category of directed augmented complexes introduced by Steiner in [9]. Following Example 3.8 of [9] and [1, §5], we have a cosimplicial object $c\colon \mathbf{\Delta} \to \mathcal{C}_{\mathrm{da}}$ from which we get a pair of adjoint functors

$$c\colon \widehat{\mathbf{\Delta}} \to \mathcal{C}_{\mathrm{da}} \quad \text{and} \quad N\colon \mathcal{C}_{\mathrm{da}} \to \widehat{\mathbf{\Delta}}.$$

We call *weak equivalences of* $\mathcal{C}_{\mathrm{da}}$ the morphisms whose image by N is a weak equivalence of $\widehat{\mathbf{\Delta}}$.

Theorem 3.6. *The functor $N\colon \mathcal{C}_{\mathrm{da}} \to \widehat{\mathbf{\Delta}}$ induces an equivalence at the level of the homotopy categories. In particular, the category of augmented directed complexes model homotopy types.*

Proof. By Theorem 2.4, it is enough to show that $c\colon \mathbf{\Delta} \to \mathcal{C}_{\mathrm{da}}$, the cosimplicial object in $\mathcal{C}_{\mathrm{da}}$, satisfies condition (e). This follows by Theorem 9.5 and Corollary 10.11 of [1], see also Remark 10.14 of [1]. □

References

[1] D. Ara and G. Maltsiniotis. *The homotopy type of the ∞-category associated to a simplicial complex*. 2015. arXiv:1503.02720.

[2] D. Ara and G. Maltsiniotis. "Vers une structure de catégorie de modèles à la Thomason sur la catégorie des n-catégories strictes". *Adv. Math.* 259 (2014), pp. 557–654.

[3] J. Chiche. "Théories homotopiques des 2-catégories". *Cah. Topol. Géom. Différ. Catég.* 56.1 (2015), pp. 15–75.

[4] R. Fritsch and D. M. Latch. "Homotopy inverses for nerve". *Math Z* 177.2 (1981), pp. 147–179.

[5] A. Grothendieck. "Techniques de construction et théorèmes d'existence en géométrie algébrique III : préschémas quotients". *Séminaire Bourbaki* 6 (1960), pp. 99–118.

[6] M. L. del Hoyo. "Espacios clasificantes de categorías fibradas". Ph.D. thesis. Universidad de Buenos Aires, 2009.

[7] L. Illusie. *Complexe Cotangent et Deformations I & II*. Vol. 239 and 238. Springer-Verlag, 1972.

[8] D. M. Kan. "On c. s. s. complexes". *Amer. J. Math.* 79 (1957), pp. 449–476.

[9] R. Steiner. "Omega-categories and chain complexes". *Homology, Homotopy and Applications* 6.1 (2004), pp. 175–200.

[10] R. Street. "The algebra of oriented simplexes". *Journal of Pure and Applied Algebra* 49.3 (1987), pp. 283–335.

[11] R. W. Thomason. "Cat as a closed model category". *Cahiers de topologie et géométrie différentielle catégoriques* 21.3 (1980), pp. 305–324.

Aix Marseille Univ, CNRS, Centrale Marseille, I2M, Marseille, France
E-mail address: andrea.gagna@univ-amu.fr